

**SEVERE COLD AND FLU DAYTIME/NIGHTTIME- acetaminophen,  
dextromethorphan hbr, doxylamine succinate, guaifenesin, phenylephrine hcl  
Target Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Target 44-640677-08**

***Active ingredients (in each caplet) (Daytime Severe Cold & Flu)***

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Guaifenesin 200 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Expectorant  
Nasal decongestant

***Active ingredients (in each caplet) (Nighttime Severe Cold & Flu)***

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Doxylamine succinate 6.25 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Antihistamine  
Nasal decongestant

***Uses***

- temporarily relieves common cold and flu symptoms:
  - minor aches and pains
  - nasal congestion
  - headache
  - fever
  - sore throat
  - sinus congestion and pressure

- cough due to minor throat and bronchial irritation
- runny nose and sneezing (**Nighttime only**)
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (**Daytime only**)

## **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- 3 or more alcoholic drinks every day while using this product
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

## **Ask a doctor before use if you have**

- liver disease
- diabetes
- high blood pressure
- glaucoma (**Nighttime only**)
- thyroid disease
- heart disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema (**Daytime only**)

- difficulty in urination due to enlargement of the prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema (**Nighttime only**)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Nighttime only**)

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children (**Nighttime only**)
- marked drowsiness may occur (**Nighttime only**)
- avoid alcoholic beverages (**Nighttime only**)
- be careful when driving a motor vehicle or operating machinery (**Nighttime only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Nighttime only**)

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Read each section carefully. Do not take DAYTIME and NIGHTTIME products at the same time.**

**Directions**

- **do not take more than directed**
- do not take more than 8 caplets of Daytime and Nighttime products in any 24-hour period
- adults and children 12 years and over: take 2 caplets with water every 4 hours
- children under 12 years: ask a doctor

**Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

- see end flap for expiration date and lot number

**Inactive ingredients (Daytime only)**

corn starch, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Inactive ingredients (Nighttime only)**

black iron oxide, corn starch, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments?**

**Call 1-800-910-6874**

**Principal Display Panel**

Compare to active ingredients in <b>Vicks® DayQuil® Severe Cold &amp; Flu* MAX STRENGTH daytime severe cold and flu acetaminophen</b> (pain reliever / fever reducer) dextromethorphan HBr (cough suppressant) guaifenesin (expectorant) phenylephrine HCl (nasal decongestant) aches, fever and sore throat chest congestion and cough nasal congestion <b>up &amp; up™ ACTUAL SIZE 16 CAPLETS</b>	
	NDC 11673-649-08 Compare to active ingredients in <b>Vicks® NyQuil® Severe Cold &amp; Flu* MAX STRENGTH nighttime severe cold and flu acetaminophen</b> (pain reliever / fever reducer) dextromethorphan HBr (cough suppressant) doxylamine succinate (antihistamine) phenylephrine HCl (nasal decongestant) aches, fever and sore throat nasal congestion and cough sneezing and runny nose <b>up &amp; up™ ACTUAL SIZE 8 CAPLETS</b>

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS**

**OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR  
SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® Severe Cold & Flu and Vicks® NyQuil® Severe Cold & Flu.

50844          REV0519A64067708

Distributed by Target Corporation

Minneapolis, MN 55403

©2019 Target Brands, Inc.

### Drug Facts

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Active Ingredients (in each caplet)**

- Acetaminophen 325 mg. Pain reliever/fever reducer
- Dextromethorphan HBr 10 mg. Cough suppressant
- Phenylephrine HCl 5 mg. Nasal decongestant

**Purpose**

Relieves pain, reduces fever, suppresses cough, and relieves nasal congestion.

**Uses**

- Temporarily relieves common cold and flu symptoms.
- Nasal congestion and pressure
- Sore throat
- Minor aches and pains
- Headache

**Warnings**

- Temporarily relieves common cold and flu symptoms.
- Nasal congestion and pressure
- Sore throat
- Minor aches and pains
- Headache

**Directions**

Take 1 caplet 4 times a day while using this product.

**Other Information**

- Keep out of reach of children.
- Do not take more than directed.
- Do not take more than directed.

### Drug Facts

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Active Ingredients (in each caplet)**

- Acetaminophen 325 mg. Pain reliever/fever reducer
- Dextromethorphan HBr 10 mg. Cough suppressant
- Phenylephrine HCl 5 mg. Nasal decongestant

**Purpose**

Relieves pain, reduces fever, suppresses cough, and relieves nasal congestion.

**Uses**

- Temporarily relieves common cold and flu symptoms.
- Nasal congestion and pressure
- Sore throat
- Minor aches and pains
- Headache

**Warnings**

- Temporarily relieves common cold and flu symptoms.
- Nasal congestion and pressure
- Sore throat
- Minor aches and pains
- Headache

**Directions**

Take 1 caplet 4 times a day while using this product.

**Other Information**

- Keep out of reach of children.
- Do not take more than directed.
- Do not take more than directed.

### daytime severe cold and flu acetaminophen

(pain reliever/fever reducer)  
dextromethorphan HBr (cough suppressant)  
guaifenesin (expectorant)  
phenylephrine HCl (nasal decongestant)

**up&up**

### nighttime severe cold and flu acetaminophen

(pain reliever/fever reducer)  
dextromethorphan HBr (cough suppressant)  
doxylamine succinate (antihistamine)  
phenylephrine HCl (nasal decongestant)

**up&up**

### daytime severe cold and flu acetaminophen

(pain reliever/fever reducer)  
dextromethorphan HBr (cough suppressant)  
guaifenesin (expectorant)  
phenylephrine HCl (nasal decongestant)

**up&up**

16 CAPLETS

### nighttime severe cold and flu acetaminophen

(pain reliever/fever reducer)  
dextromethorphan HBr (cough suppressant)  
doxylamine succinate (antihistamine)  
phenylephrine HCl (nasal decongestant)

**up&up**

8 CAPLETS

### Drug Facts (continued)

**Directions**

Read each section carefully. Do not take DAYTIME and NIGHTTIME products at the same time.

**Warnings**

- Do not take more than directed.
- Do not take more than directed.

**Other Information**

- Keep out of reach of children.
- Do not take more than directed.

### Drug Facts (continued)

**Directions**

Read each section carefully. Do not take DAYTIME and NIGHTTIME products at the same time.

**Warnings**

- Do not take more than directed.
- Do not take more than directed.

**Other Information**

- Keep out of reach of children.
- Do not take more than directed.

Target 44-640677

No print/No varnish Lot & Exp date

094 14 0107 R01  
IDC-000274-01-028

3 50844 64908 9

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® Severe Cold & Flu and Vicks® NyQuil® Severe Cold & Flu.

50844 REVO519464067708

Distributed by Target Corporation  
Minneapolis, MN 55403  
©2019 Target Brands, Inc.

### Drug Facts (continued)

**Other Information**

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

**Inactive Ingredients (Daytime only)**

acetaminophen, dextromethorphan HBr, guaifenesin, phenylephrine HCl, sodium starch glycolate, stearic acid, titanium dioxide

**Inactive Ingredients (Nighttime only)**

acetaminophen, dextromethorphan HBr, doxylamine succinate, phenylephrine HCl, sodium starch glycolate, stearic acid, titanium dioxide

**Questions?** Call 1-800-910-6374

B-1920-640677-08  
REVO519464067708

## SEVERE COLD AND FLU DAYTIME/NIGHTTIME

acetaminophen, dextromethorphan hbr, doxylamine succinate, guaifenesin, phenylephrine hcl kit

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-649
--------------	----------------	--------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-649-08	1 in 1 PACKAGE, COMBINATION; Type 0: Not a Combination Product	08/01/2015	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 BLISTER PACK	16
Part 2	1 BLISTER PACK	8

## Part 1 of 2

## SEVERE COLD AND FLU DAYTIME

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

### Product Information

Route of Administration	ORAL
-------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	

<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	44;640
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	02/27/2014	

### Part 2 of 2

#### SEVERE COLD AND FLU NIGHTTIME

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl tablet, film coated

### Product Information

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: QD3BT10XU1)	DEXTROMETHORPHAN	



<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9DZRT19KTH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
<b>FERROSOFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	44;677
<b>Contains</b>			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/01/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/01/2015	

**Labeler** - Target Corporation (006961700)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11673-649)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11673-649)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	pack(11673-649)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11673-649)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	pack(11673-649)

Revised: 11/2021

Target Corporation